

Anti-Mouse CD117 (c-Kit) SAFIRE Purified

Catalogue Number : 19112-25

RUO: For Research Use Only. Not for use in diagnostic procedures.

Product Information

Clone: ACK2

Format/Conjugate: SAFIRE Purified

Concentration: 2 mg/mL

Reactivity: Mouse

Laser: Not Applicable

Peak Emission: Not Applicable

Peak Excitation: Not Applicable

Filter: Not Applicable

Brightness (1=dim,5=brightest): Not Applicable

Isotype: Rat IgG2b, kappa

Formulation: Phosphate-buffered aqueous solution, pH7.2.

Storage: Product should be kept at 2-8°C and protected from prolonged exposure to light.

Applications: FC, FA, IHC, ICC, WB

Description

The ACK2 monoclonal antibody specifically reacts with mouse CD117 (c-Kit receptor), a 145 kDa transmembrane tyrosine-kinase receptor encoded by the Kit gene. The c-Kit receptor, also known as stem cell factor receptor, is expressed on hematopoietic progenitor cells in adult bone marrow, in progenitors of erythroid and myeloid lineages, and precursors of B and T cells. CD117 enhances the proliferation and the differentiation of the hematopoietic progenitor cells and seems to enhance the development of T cells, as the c-Kit receptor and its ligand are expressed by the thymus.

Preparation & Storage

The product should be stored undiluted at 4°C. Do not freeze. The monoclonal antibody was purified utilizing affinity chromatography. The endotoxin level is determined by LAL test to be less than 0.01 EU/µg of the protein.

Application Notes

The antibody has been analyzed for quality through the flow cytometric analysis of the relevant cell type. It is recommended that the reagent be titrated for optimal performance for each application.

References

- Godfrey, D. I., Kennedy, J., Mombaerts, P., Tonegawa, S., Zlotnik, A. (1994). Onset of TCR-beta gene rearrangement and role of TCR-beta expression during CD3-CD4-CD8-thymocyte differentiation. *The Journal of Immunology*,;152(10), 4783-4792.
- Godfrey, D. I., Zlotnik, A. L. B. E. R. T., Suda, T. A. K. A. S. H. I. (1992). Phenotypic and functional characterization of c-kit expression during intrathymic T cell development.;*The Journal of Immunology*,;149(7), 2281-2285.
- Feng, H., Sandlow, J. I., Sandra, A. (1998). The c-kit receptor and its possible signaling transduction pathway in mouse spermatozoa.;*Molecular reproduction and development*,;49(3), 317-326.